

Index to Subjects

Many topics are treated in numerous places throughout individual chapters and in the many different chapters. This index gives detailed citations even for relatively scattered entries. Readers seeking a more general overview of content may prefer for some topics to consult the Contents (page vii) or the Abstracts that precede each chapter.

For taxa, see especially Index to Families and Other Higher Taxa of Arthropods, page 1015; Contents, page vii; and annotated lists for particular Orders or Families.

Not indexed here, because they would have resulted in thousands of additional entries, are the following:

- Specific data for individual taxa given in annotated lists.
- Information in Abstracts.
- Aspects of general biology, as opposed to specific Yukon information, treated in the introductions for the various taxa.
- Individual place names (but for general orientation see especially pp. 13–57, and for localities see especially pp. 333–336, 341–343).

- Adaptability, of leafhoppers, 369–370
Adaptations, 984–987 (see also Cold hardiness...;
 Diapause; Life cycle; Phenology)
 arctic, alpine, boreal or northern, 8–11, 118, 366, 401,
 604, 608, 627–633, 825, 831, 851, 861, 971,
 976, 985, 987, 1005, 1008
 lack of, 244, 340, 370, 688, 828, 1005
 of mammals, 60, 64
 to Beringian conditions, 480, 550, 649–650, 833, 1003
 to food or its timing, 221, 608
 to limited resources, 984–985 (see also Resources)
 to peatlands, 46
 to postglacial conditions, 840, 843
 to saline habitats, 46, 244
 to seasonality, 985–986 (see also Life cycle;
 Phenology)
 to shallow habitats, 520
 to transient habitats, 828, 860
Additional or other species expected in Yukon, 105–
 107, 160–161, 225–226, 359, 396–400, 514–
 515, 541–543, 587–593, 627, 646, 673–674,
 712–713, 775, 893–894, 907, 971, 972
Adelphotaxon, 476, 477
Adjacent regions to Yukon, faunas of, 192, 198, 225,
 289, 291, 300, 360, 396, 409–410, 419, 514,
 616, 627, 641, 665, 820, 855, 893, 894, 908,
 975–976 (see also Alaska; Northwest
 Territories)
Adventive species, 224, 231, 237, 322
Aestivation, 224
Age of species, 8, 146, 219, 370, 372, 824, 825
Age of tundra habitats, 825, 917
Alaska, fauna of, 74, 117, 120, 122, 161, 163, 192,
 198, 225, 289, 291, 300, 336, 340, 360, 379,
 384, 396, 409–410, 419, 454, 514, 517, 518,
 522, 553, 587, 627, 628, 641, 689, 692, 712–
 713, 779, 820, 855, 856, 907, 918, 975–976
 (see also entries in annotated lists)
Alkaline pond, 172, 981
Allochthonous food, 822, 858, 859
Allochthonous fossil deposits, 941, 949
Alpine glaciers, 17, 19, 23, 54
Alpine or Arctic/alpine species or fauna, 8, 9, 10, 107–
 109, 221, 323, 326, 339, 340, 361, 366, 369,
 518, 519, 678, 716, 718, 723, 825, 849, 947,
 949, 1009 (see also Adaptations; Mountain;
 Tundra)
 origin of, 825, 854
Alpine vegetation or zone, 38–41, 44, 53, 54, 340,
 343, 467, 980
 in past, 917, 957
Alvar, 370
Analogues or lack of analogues to Pleistocene envi-
 ronments, 61–62, 64, 68, 553, 916, 947, 1002
Antennae, loss or reduction in chironomid midges,
 632–633, 979
Anthropochorous or Anthrophilic species, 401, 402,
 565, 568 (see also Introduced; Synanthropic)
Ants, and secretions of bugs, 243
 resemblance to, see Myrmecomorphy
Aposematism, in true bugs, 243, 244, 308
Aptery, see Wings, loss or reduction
Aquatic communities, and caddisflies, 791, 857, 858
 and dragonflies, 171
 and water beetles, 518–519
Aquatic habitats in Yukon, 171, 413, 492, 493, 500,
 627, 791, 861, 981–982, 984, 1008 (see also
 Lake; Pond; River; etc.)
Aquatic versus terrestrial habitats or species, 987–
 989, 1008
Arboreal species, 117, 118, 449, 455, 467
Arctic islands refugium, 220
Arctic vegetation, environment or zone, 9, 38, 39, 40,
 52, 340, 343, 467, 468, 493, 780, 791, 965, 980,
 1002 (see also Fell-field)
 in past, 326, 366, 368, 608, 825, 959, 1000, 1002
Arctic species or fauna, 9, 10, 107–109, 117, 146,
 163, 219, 230, 244, 300, 301, 320, 322, 340,
 360, 361, 363, 365, 366, 370, 383, 384, 388,
 401, 470, 471, 495, 516, 517, 518, 526, 615–
 632, 650, 716, 717, 718, 719, 723, 783, 828,
 830, 849, 861, 895, 976, 997–999, 1006, 1007,
 1009 (see also Adaptations; Alpine...; Tundra)
 origin of, 470, 825, 830, 849, 1006

- Arctic, Tertiary floras and faunas in, 951
- Arctic steppe, *see* Steppe
- Aridity, Arid habitats, 30, 48, 236, 364, 605
in Pleistocene/Beringia, 64, 65, 68, 108, 477, 480, 823
- Artemisia*-grass communities, 45, 48–50, 67, 302, 325, 369, 778 (*see also* Sage)
- Asiamerican origin, 218, 219
- Aspect, as determinant of species richness, 897
- Attenuation, *see* Diversity and latitude
- Autochthonous Beringian origin or species, 481, 606
- Autochthonous fossil deposits, 941, 945, 948, 956
- Autochthonous food, 822, 859
- Autogeny or autogenous, in black flies, 566, 604, 605, 607, 608, 985
- Barriers, *see* Dispersal, barriers to
- Basking, 984
- Bering or Beringian land bridge, 60, 66, 67, 68, 109, 110, 220, 236, 238, 365, 366, 367, 469, 790, 821, 822, 823, 824, 830, 831, 916, 917, 956, 958, 959, 1000
dispersal or failure to disperse across, 826, 829, 835, 836, 837, 838, 839, 841, 842, 843, 849, 850, 851, 852, 859, 958
habitats or conditions of, 60, 61, 63–68, 365, 369, 823, 832, 833, 859, 916, 953, 958, 959, 1003
- Bering Strait or Sea, 23, 28, 64, 109, 220, 366, 609, 649, 779, 790, 850, 966, 1000, 1008
as a barrier, 110, 721, 823
crossing of, 361, 823, 1003, 1008
not a separator or species similar, 8, 219, 372, 456, 656, 720, 721, 1004 (*see also* Palaearctic region, connections...)
separator or species different, 109, 110
- Beringia, 7, 8, 54, 60, 61, 370, 454, 469, 481, 550, 608, 646, 679, 720, 916, 1001 (*see also* Endemic species; Glaciation; Ranges; Refugia; and entries below)
and antennal reduction, 632, 633
dispersal or movement, across or via, 191, 220, 238, 323, 1003
into, 325, 365, 779, 822, 839, 843, 1006
out of, 10, 471, 481, 822, 830, 833, 836, 843, 850, 1005
out of, lack of movement, 652, 841, 842, 849
early history of, 858–859, 958
extent or limits of, 8, 408–409, 455, 469, 565, 605, 725, 790, 822–823, 916, 958, 966, 989, 1001
past fauna of, *see* Fossils
past habitats or conditions in, 23, 49–50, 59–72, 108, 109, 198, 324, 325, 326, 367, 368, 448, 470, 471, 477–481, 550–553, 605–606, 608, 650, 718, 719, 779, 821, 823, 851, 852, 859–860, 916, 947–951, 979, 1001–1002, 1003, 1009
during Tertiary, 951–955, 958
past vertebrate fauna of, 60, 61, 67, 605–606, 916, 948
survival or evolution in, 108, 145, 236, 325, 470, 475, 550, 606, 719, 909, 978, 1002 (*see also* Refugia)
- Beringian centrant species, in black flies, 606, 607, 608
in ground beetles, 470–471, 477, 479, 480, 481
- Beringian emigrant species, in black flies, 606, 607, 608
in ground beetles, 474, 475
- Beringian endemic species, *see* Endemic species
- Beringian fauna and Yukon project, 1, 2
- Beringian habitats, *see* Beringia, past habitats...
- Beringian immigrant species, in black flies, 606
in ground beetles, 474, 475, 476
- Beringian isolates, 412, 415, 416, 476, 551
- Beringian lineages, 372–373
- Beringian paradox, 841, 849
- Beringian refugium, *see* Refugia or refugium, Beringian
- Beringian resident species, in black flies, 606, 607, 608
in ground beetles, 474, 475
- Bicentric, *see* Range, bicentric
- Biennialism, in butterflies and moths, 783, 986
- Biogeographic categories or patterns, *see* Ranges
- Biogeography, *see especially* Range types
- Biological filter or filter, 365, 958
- Biological Survey Foundation, 4
- Biological Survey (of Canada), 1–5, 7, 903
- Birds, and insects, 569 *et seq.*, 604–608, 903, 909
- Blood feeding or bloodsucking, 565, 568, 597, 604, 605, 608, 985
- Bog, 45–46, 63, 109, 119, 173, 301, 378, 447, 491, 500, 627, 843, 981 (*see also* Peatland)
- Bog tundra, 119, 130
- Boreal forest, vegetation or zone, 7, 8, 14, 38, 43–44, 48, 53, 54, 64, 344, 363, 467, 493, 679, 949, 965, 976, 980
in past, 68, 109, 917, 953, 954, 955, 957, 958
- Boreal or boreal-forest species, 9, 10, 11, 107–110, 119, 130, 144, 145, 146, 147, 175–176, 359, 361, 364–365, 366, 370, 383, 396, 403, 477, 516–520, 544, 545, 549, 550, 551, 640, 646, 654–656, 674–679, 719, 776–783, 827, 828, 897, 909, 970, 997–999, 1005, 1006, 1007, 1008, 1009 (*see also* Adaptations; Ranges, Boreal)
as predominant in Yukon, 164, 171, 192–193, 361, 383, 413, 515, 553, 779, 988, 997
representation in collections, 339, 725, 970, 997
- Boreomontane species, 130, 146, 189, 402, 471, 544–545, 549, 550, 551, 552, 553, 640, 677, 679, 721, 726, 776–777, 779, 780, 781, 782, 825, 997, 998, 999 (*see also* Ranges, Boreomontane)
- Bottleneck for survival, 950, 1009
- Brachyptery, *see* Wings, reduction or loss
- British Columbia, fauna of, 122, 192, 194, 197, 198, 199, 203, 225, 289–290, 291, 300, 302, 360, 366, 379, 396, 514, 518, 522, 587, 646, 641, 854–855, 856, 971, 975–976
- Canadian Arctic Expedition, 244, 447
- Canyons, 16
- Carrion, 675, 943, 948
- Cascadia, 220
- Cave, 678

- Bluefish fossil site, 60, 916, 948
- Checklists of species, 122–135, 160–161, 163, 177, 205–206, 235, 291–299, 449–454, 528–529, 566–567, 628–631, 640–641, 675–676, 714–715, 826–827, 830–831, 841, 850, 904 (see also the annotated lists)
- Clade, 677, 679
- Classification, of aquatic habitats, 152, 162, 164 (see also Aquatic habitats)
- of arthropods, see checklists and annotated lists of ranges, see Range types; Ranges
- of soils, 35
- Cleptoparasites, 689, 898–899, 984
- Climate, and aquatic communities, 791, 859, 861
- and changes during past, 8, 37, 59, 62, 65, 66, 68, 109, 326, 368, 459, 460, 475, 820–822, 824, 825, 897, 915–918, 946, 947, 948, 949, 1000, 1001, 1002, 1006, 1008 (see also Palaeoclimate)
- in Tertiary, 953–957, 957–958
- response to, 361, 363, 721, 946, 957, 949, 950, 1006–1007
- and diversity, 220, 976 (see also Composition...)
- and life cycles, 118 (see also Life cycle)
- and ranges, 460 (see also Ranges)
- of Yukon, 9, 10, 26–35, 38, 44, 50, 171, 896 (see also Summer temperature; Temperature; Wind)
- climatic regions or zones, 30–35
- climatic regions or zones, and leafhoppers, 338, 340–344, 346, 361–363
- models, 61, 64, 65, 68
- Coarse Particulate Organic Matter (CPOM), in streams, 220, 221, 982
- Coastal Plain, Arctic or Yukon, 8, 14, 23, 24, 38, 45, 50, 51, 52, 53, 119, 136, 221, 227, 338, 384, 664, 725, 792, 835, 895, 897, 945, 976, 981, 1009
- Coastal species, 119, 366, 369
- Cold hardiness, cold tolerance or cold adaptation, 118, 219, 220, 221, 225, 235, 236, 237, 322, 477, 608, 627, 790, 821, 858, 859, 986–987, 1003, 1008, 1009, 1010
- Collecting in the Yukon, 1–4, 968–970
- for aculeate wasps, 870
- for anthomyiid flies, 692
- for ants, 902–903
- for aphids, 388, 389, 401
- for black flies, 568–569
- for butterflies and moths, 724–725, 775
- for caddisflies, 791
- for chironomid midges, 615–616
- for delphacid planthoppers, 379
- for dragonflies, 171
- for ground beetles, 447–448
- for leafhoppers, 338–340
- for mayflies, 152, 153, 161
- for oribatid mites, 117, 118
- for orthopterans, 226
- for robber flies, 638–639
- for sphaerocerid flies, 664–665
- for spiders, 75
- for stoneflies, 203
- for true bugs, 3–4, 244–245
- for water beetles, 492
- needs for or inadequacy or deficiencies of, 48, 110, 119, 142, 145, 147, 161, 164, 226, 339, 360, 365, 388, 401, 517, 594, 609, 615, 627, 639, 679, 720, 725, 775, 776, 782, 896, 899, 969–970, 972–974, 975, 997, 1010
- Collecting sites in the Yukon, located or mapped, 154, 178, 246, 333–336, 341–343, 649 (see also localities in annotated lists and appendixes)
- of fossils, 465–466, 913
- Colonization or postglacial colonization or recolonization, 164, 193, 218, 364, 401, 471, 604, 654, 719, 721, 835, 837, 843, 846, 859, 860, 897, 982 (see also Invasion)
- Colour, variation in, 812, 836, 837, 839, 840, 850, 978 (see also Aposematism)
- Comparison of faunas, see Fauna
- Competition or competitors/competitive, 220, 221, 840–841, 849, 851–852, 982, 984, 1008, 1010
- Composition or diversity of the Yukon fauna, 971–978, 987–989, 1007–1009 (see also Diversity and latitude...)
- compared to elsewhere, see Fauna
- in aculeate wasps, 894–897
- in anthomyiid flies, 689
- in ants, 907–909
- in aphids, 401
- in beetles, 409–410, 416
- in black flies, 593–595
- in butterflies and moths, 775–776
- in caddisflies, 793, 854, 855, 858
- in chironomid midges, 626–632
- in delphacid planthoppers, 379
- in dragonflies, 176–177
- in ground beetles, 454–455
- in leafhoppers, 360–361
- in mayflies, 161–164
- in oribatid mites, 145
- in orthopterans, 225
- in robber flies, 646
- in sphaerocerid flies, 664–665
- in spiders, 107–108
- in stoneflies, 203
- in true bugs, 289, 300
- in water beetles, 518–519
- in weevils, 543
- Coniferous forest, and soils, 37
- dark and light, 917, 959
- species associated with, 109, 302, 324, 325, 656 (see also Boreal; Boreomontane)
- Cooperation or cooperative, 1, 2, 4, 5, 7, 482
- Cordillera, aquatic habitats, 604, 980, 989, 1003, 1005
- ecozones, 52, 53, 245, 300, 302, 307, 870, 896
- movement along or route, 10, 1003, 1006–1007, 1009
- relicts in or refuge, 10, 1009 (see also Refugia...; Relict)
- species in, 107, 108, 109, 110, 143, 146, 162, 164, 173, 175, 176, 189, 192, 193, 194, 199, 640,

- 653, 654, 678, 716, 717, 989 (and see especially Ranges, Cordilleran...; see also Mountain) sister taxa of, 677
uplift of, 65, 825
- Cordilleran ice, ice-sheet or glacier, 9, 23, 60, 65, 652, 790, 813, 821, 822, 824, 830, 840, 848, 913, 1000–1001, 1003, 1006
- Cordilleran physiographic region, 15, 38
- Corridor, see Dispersal
- Cryoturbation, 36–37
- Cul-de-sac, Yukon as, 916, 948
- Cytospecies or cytologically recognized species, 565, 569, 593, 606
- Data, strengths or weaknesses of, summary, 968–970 (see also Collecting, needs for...)
- Deciduous forest, water beetles associated with, 493, 499, 517
- Decomposition, or limitation of, 116–117, 983, 984
- Deglaciation, response to, 8, 9, 68, 365, 471, 551, 604, 606, 779, 780, 828, 835, 837, 840, 843, 846, 897, 966, 982, 1003, 1006, 1007 (see also Glaciers, retreat of; recession of)
- Desiccation, of habitats, 145, 860, 861
resistance to, 118, 828, 860
- Detritivores or Detritivory, 204, 858, 859, 861, 988
- Detritus, in streams or ponds, 221, 827, 838, 854, 858–859, 981, 983
with fossils, 944, 947
- Development, rate of or time for, 118, 171, 195, 196, 204, 783, 858, 860, 983, 984, 986
- Diapause, 204, 221, 224, 783, 828, 860, 986, 987
and latitude, 860, 986
- Differentiation, see Evolution; Isolation; Speciation; Variation
- Disjunct, see Ranges
- Dispersal, abilities or lack of abilities for, 164, 198, 219, 304, 306, 364, 365, 401, 460, 468, 470, 474, 481, 604, 719, 721, 823, 828, 987, 1003, 1006
across Beringia/land bridge, 323, 324, 372, 823, 824, 830, 831, 832, 833, 837, 838, 839, 841, 842, 843, 849, 853, 854, 859
barriers to, 459, 632, 958, 1003
 Bering Strait, 110, 721, 823
 ice sheets, 652, 1003
 in Cenozoic, 821
 mountains, 10, 11, 162, 171, 194, 364, 953, 957, 1003
 ocean, 632, 721, 821, 823
corridor for, across Beringia/land bridge, 23, 823, 835, 916, 958–959
 ice-free between ice sheets, 23, 822, 830, 917, 1001
 river systems as, 220
eastward out of Beringia, 323, 474, 822, 839, 850, 851, 1006
from south, 326, 843
into Beringia, 477, 551, 823
into Greenland, 823, 828, 843, 846
northward, 324, 366, 551, 832
of anthomyiid flies, 719, 721
of aphids, 401
of black flies, 604, 607
of caddisflies, 821–823, 828–837, 843, 846, 848–854, 860
of dragonflies, 198, 823
of ground beetles, 460, 468, 470, 474, 477, 481
of leafhoppers, 339, 364–369, 969
of mayflies, 164
of orthopterans, 235–237
of true bugs, 304–306, 322–326
of weevils, 551–552
post-Pleistocene, 322, 323, 324, 326
southward, 604, 826, 833
westward, 365, 412, 552
(see also Bering...land bridge, dispersal...; Migration; Movement)
- Distribution or Distributional categories, elements or patterns, see Ranges (and see especially data in annotated lists for individual species)
- Distribution patterns within the Yukon, 136–141, 188–196, 234, 235, 361–363, 384, 517–520, 640–654, 976
- Distributions, restricted, see Endemic; Restricted
- Disturbance, of soil microhabitats, 119 (see also Cryoturbation)
as habitat feature, 44, 302, 466, 984
- Diurnal activity, 775, 985
- Divergence, morphological or structural, 552, 824, 835, 846
rate of, 825
time of, 326, 1004
- Diversity and area, 519–520, 522, 971
- Diversity and latitude—decline or attenuation, 976
in aculeate wasps, 896
in beetles, 414
in caddisflies, 827, 854–859
in dragonflies, 196–199
in leafhoppers, 339, 361–363
in true bugs, 300–301
in water beetles, 518–520
in weevils, 526
- Diversity, depleted by cooling and glaciation, 957
- Diversity or richness in Yukon, see Composition
- DNA, studies using, 289, 470, 606
- Dormancy, 204
- Drainage systems, 17, 23–26, 54, 161, 302, 366, 368, 791
changes in, 822, 917, 1001, 1006
connections between, 162, 164, 1003
- Dry habitats, and species associated with, 115, 302, 416–417, 449, 466, 467, 475, 476, 480, 523, 637, 654, 655, 678, 717, 723, 897 (see also Arid; *Artemisia*-grass; Dune; Steppe)
in past, 365, 650, 860, 947, 948, 949, 951, 1002, 1003
- Dry tundra, and species associated with, 8, 119, 130, 142, 144, 145, 146, 147, 301, 324, 415, 445, 468, 471, 523, 544, 545, 549, 550, 553, 674–675, 678, 718, 719, 723, 776–782, 981, 1002, 1006
in past in Beringia, 61, 64, 369, 553, 1002, 1009

- Dunes or sand, 9, 48, 65, 130, 142, 370, 413, 449, 466, 467, 474, 479, 480, 655, 691, 718, 719, 776–778, 780–782, 784, 998, 999
- Dung, 415, 675, 904, 983
- Ecogeographic region, 2, 50–51, 53, 54, 136, 153, 154, 161, 227, 234, 235, 379, 493, 527, 665, 791–793, 855
- Ecological correlations with faunal composition or diversity, 987–989
- Ecological generalist or generalist, 302, 413, 467, 468, 469, 474, 475, 476, 477, 479, 480, 547, 678, 831, 841, 1008 (see also Hostplant)
- Ecological niche, 234, 791, 854, 857
vacant, 823
- Ecological relationships or considerations, overview, 979–989 (see also accounts for individual taxa; and more detailed index entries such as Habitat)
- Ecological testing ground, Yukon as, 791
- Ecoregion, 14, 50, 52–54, 178, 246, 247, 248, 301, 460, 641, 870, 895, 896, 897
- Ecozone, 14, 50, 52–54, 245, 246, 300–301, 302, 307, 870, 896
- Eggs, number of in black flies, 608
resistant to desiccation in caddisflies, 828, 869
overwintering as, 196, 220–221, 224, 378, 986
- Elevation, as determinant of aculeate species richness, 897
species at high, 220, 724, 782, 783 (see also Adaptations...; Mountain)
- Emergence, time of, 340, 627, 970, 985, 986 (see also Biennialism; Phenology)
- Emigrants, see Beringian emigrants
- Endemic genera, in North America, 218
- Endemic species, and habitats, 415, 474, 480, 552–553, 652, 719, 780–781, 1002
in Alaska, 469
in Beringia, 8, 9, 205, 218, 219, 300, 321, 326, 338, 339, 360, 361, 363, 364, 365, 366, 369, 370, 372, 373, 412, 415–416, 470, 471, 474–476, 481, 552, 597, 604, 606, 674, 678, 689, 715, 718, 719, 725, 778, 779–782, 849, 850, 851, 1004, 1005, 1008, 1009 (see also Restricted species; and see Ranges, Beringian)
in mountains, 10, 370, 597 (see also Mountain; Refugia)
in northwestern North America, 369, 677
in single life zone, 363
in Yukon, 49, 145, 223, 224, 238, 341, 368, 377, 454, 781–782
- Endemism, areas of, 341
- Environment of the Yukon, current, 13–57
- Equilibrium, or lack of, between fauna and climate, 10, 66, 68, 1001, 1007
- Evidence, to explain faunal composition and history, 999–1000, 1004–1005
- Evolution of species, in leafhoppers, 369–373
in orthopterans, 234–238
rate of, 825, 914, 949, 954
(see also Isolation; Speciation; Variation; and other specific topics)
- Extinct insect species, or extinction, 369, 370, 608, 953, 954, 957, 958, 823, 825, 840, 851, 852, 915, 918, 951, 955
- Eyes, ellipsoid or reduced, 9, 852, 942, 943, 987
- Fauna of Yukon, compared to elsewhere, 161–162, 196–198, 289–300, 360, 409–410, 454, 518–520, 594–595, 605, 627, 775, 854–856, 894, 971–976 (see also Alaska, fauna of; and see similar entries)
- Faunal elements, see Ranges
- Fecundity, 118, 306
- Feeding or trophic guild, 303, 318, 322, 325, 856, 858
- Feeding habits, see Food habits
- Fell-field, 62, 119, 365, 416, 477, 552, 948, 1002
- Fen, 45, 46, 172, 173, 174, 175, 301, 675, 981
- Fieldwork, see Collecting
- Filter, see Biological filter
- Filter feeding, decline in Yukon, 856–857, 983, 984
loss of habit in black flies, 608, 983
- Fine Particulate Organic Matter (FPOM), in streams, 856, 857
- Fire, 44, 68, 324, 915, 980
- Flightlessness, see Wings, loss or reduction
- Food or Feeding habits, 981, 982–985 (see also Hostplant; and for specific information see annotated lists)
of aculeate wasps, 898, 899
of aphids, 401
of beetles, 413–415, 416
of black flies, 566, 604–605, 607, 608
of butterflies and moths, 779
of caddisflies, 822, 827, 836, 856–859
of leafhoppers, 339, 361, 366, 369, 372
of oribatid mites, 116, 118, 119, 142, 144, 145
of stoneflies, 221
of true bugs, 302–304, 318, 325
of weevils, 546–547, 552
- Food web, in aquatic habitats, 204, 217, 982
- Foodplants, see Hostplants
- Forest, as habitat for ground beetles, 467, 469, 475, 476, 479, 480, 481
distribution of arctic species in, 830
movement of tree species in past, 68, 363, 364, 915, 945, 946
(see also Boreal forest; Coniferous forest; Deciduous forest; Taiga; see also Ranges such as boreal)
- Fossils, arthropod or insect, 63, 64, 68, 143, 236, 323, 360, 366, 454, 459–478, 527, 541, 551, 552, 606, 691, 825, 909, 911–962, 968, 976, 1001, 1002, 1006
abundance in Yukon, 912, 917
from Greenland, 931, 945, 956–957
Holocene, 912, 919–923, 945–946
illustrated, 942, 950
Interglacial, 471, 949, 950–951, 1001
level of identification of, 914–915
limitation or inadequacy of, 217, 236, 238, 323, 606, 608
mid to late Pleistocene, 912, 924–930, 945–951
Modern or extant, 919–923, 944–945

- Pliocene, 143, 552, 941, 953–955, 956
 Tertiary (and early Pleistocene), 912, 917,
 931–940, 951–957
 pollen or plant, 60–65, 68, 480, 917, 945, 946, 947,
 949, 951, 953–956, 959
 preservation of, 914, 918, 941, 949
 sites for, 60, 465–466, 913 et seq.
 vertebrate, 60, 61, 67, 606, 916, 948, 956
 Founder effect, 833
 Freezing or frozen, in aquatic habitats, 24, 220, 857,
 858, 859, 945, 981, 986–987 (see also Cold
 hardness...; Ice)
 Fungi, and decomposition in aquatic habitats, 827,
 857, 983
 Fungivores or fungivory, 142, 415, 525, 546, 664,
 674, 675, 676, 677, 983, 978
 Gene flow, restricted, 721, 833, 837
 Generalist, see Ecological generalist
 Genetic systems, in mites, 118
 Geographical patterns, see Ranges
 Geology, 60, 999
 of Yukon, 14, 19
 Gills of stoneflies, and adaptation, 220, 987
 Glaciation or Glaciated area, 10, 14, 18, 22–25, 52,
 53, 59–60, 65, 162, 324, 326, 363, 366, 455,
 517, 544, 551, 596, 597, 822, 823, 851, 915–
 917, 957, 958, 997, 999–1001 (see also Re-
 fugia; Unglaciated)
 Glaciations, number of, 23, 1000, 1001
 timing of, 915–916
 Glaciers, current, 17, 19, 35, 39, 54
 Glacier-fed or glacial rivers or streams, 173, 823, 857
 Glaciers, recession of, responses to, 219, 790, 821,
 822, 826, 828, 831, 837, 839, 843, 849, 851,
 857, 1001 (see also Deglaciation; Glaciers,
 retreat...)
 retreat of, responses to, 10, 193, 323, 364, 833, 835,
 836, 839, 843, 851, 854, 858, 897, 1001, 1006
 (see also Deglaciation; Glaciers, recession...)
 Grasses and sedges, association with, 302, 303, 306,
 308, 318, 322, 323, 339, 361, 378, 384, 981,
 983 (see also Sedge)
 Grassland adaptations, in mammals, 60
 Grassland species, 1, 3, 48, 49, 146, 325–326, 339,
 365, 366, 369, 373, 493, 495, 516, 517, 656,
 827, 830, 998, 999, 1002, 1003
 Grassland soils, 38
 Grassland, vegetation, 48, 49, 897, 980 (see also
 Artemisia-grass...)
 as barrier, 205
 in past, 60, 62, 63, 64, 66, 67, 68, 69, 367–369, 370,
 477, 480, 1002
 Greenland, caddisflies in, 821, 823, 828–830, 833,
 837, 843, 845, 846
 other species in, 74, 108, 224, 244, 365, 627, 628, 631
 fossils from, 931, 945, 956–957
 Growing season, 195, 344, 860, 986 (see also
 Season...)
 Growth, rate or duration of, 196, 220, 639, 860, 986
 Habitat, and wing development, see Wings...and
 habitat
 Habitat requirements, specific, 63, 664
 Habitats in Yukon, 38–50 (see also Aquatic habitats;
 and individual habitats, e.g. Dry tundra;
 Meadow)
 heterogeneity or variety of, 9, 172, 604, 980
 mosaic, 44, 597, 980–982, 1008
 mosaic during Pleistocene, 49, 62, 63, 64, 68, 325,
 365, 368, 479, 481, 552, 605, 1002
 photographs of, iii, 20, 21, 27, 40, 41, 42, 43, 47, 173,
 174, 175, 176, 655, 656, 657
 Habitats, aquatic, see Aquatic habitats
 of Beringian species, see Beringia, past habitats or
 conditions in
 of insects, change of with latitude, 117, 835, 860,
 980, 982
 notable, 45–50
 stability or persistence of, 304, 306, 307, 339, 378,
 384, 468, 480, 481, 852, 678, 857–858, 949,
 979, 981, 982, 985, 988, 1008
 terrestrial, see Terrestrial habitats
 Habitats or Habitat associations in Yukon, 980–982,
 985, 986, 987–989, 997, 1006, 1008
 of aculeate wasps, 896, 897
 of anthomyiid flies, 719
 of ants, 904
 of aphids, 389
 of beetles, 411, 412, 413, 416
 of butterflies and moths, 776–782
 of caddisflies, 827, 830, 835, 837, 849, 851, 854–
 859, 860, 861
 of chironomid midges, 615–616, 627, 632
 of delphacid planthoppers, 384
 of dragonflies, 171–173, 198
 of ground beetles, 449, 460, 464–468, 493
 of oribatid mites, 117–118, 119, 142–146, 130
 of robber flies, 639, 655
 of sphaerocerid flies, 665, 674–679
 of spiders, 108–110
 of stoneflies, 220, 221
 of true bugs, 300–302, 308, 318
 of water beetles, 492, 500, 515–520
 of weevils, 544–546, 550, 551, 552, 553
 Herbivores or Phytophages, 119, 144, 221, 243, 302–
 304, 307, 308, 318, 323, 412, 413–415, 416,
 526, 982–983, 984, 988, 989, 1008 (see also
 Hostplants)
 Highway, see Road
 Hindwing development, see Wings
 History of collecting, see Collecting
 Holarctic species, see Ranges
 Host preferences, of aculeates, 898–899
 Hostplant alternation, in aphids, 401, 403
 Hostplant defences, as causing biennialism, 783, 986
 Hostplants or foodplants in Yukon, 982–983, 1002
 (see also annotated lists for hostplants or food-
 plants of individual species)
 of aphids, 401, 403
 of anthomyiid fly, 689, 718–719

- of beetles, 413–415
- of butterflies and moths, 776, 779
- of leafhoppers, 339, 340
- of true bugs, 302–304, 308, 322–324
- of weevils, 546–547, 549, 552, 553
- Hummocks, 36, 41, 43
- Hybridization or hybrids, 9, 192, 198, 199, 361, 606, 1005
- Hypopygium, enlarged, 632, 979
 - fossil, 943, 949
- Hyporheic or hyporheal, 220, 857, 981, 986, 1008
- Hypsithermal period, 366, 367, 897, 915, 916, 1001, 1002
- Ice, and aquatic habitats, 172, 220, 221, 857, 858, 859, 860, 983
- Ice field, see Glaciation; Glaciers
- Ice-free corridor, see Dispersal, corridors for, ice free
- Ice-front conditions or environments, 369, 370
- Ice-sheet, see especially Cordilleran ice; Laurentide ice; see also Refugia; Dispersal, barriers to; and other specific topics
- Immigrants, postglacial, see Beringian immigrants
- Information, see Data
- Insolation, 65, 861, 945, 980
- Interbreeding, 238 (see also Hybridization)
- Interglacial, 9, 10, 60, 821, 822, 917
 - conditions during, 63, 66, 366, 477, 480, 915, 1001, 1002
 - fossils, see Fossils
 - Holocene as an, 945, 1001
 - movement of organisms during, 63, 66, 471, 830, 915
- Interstadial, 61, 366, 946, 949, 1000
- Intermontane fauna, 360, 366, 373, 493
- Introduced species, 322, 326, 339, 544, 547, 548, 720, 985, 1007
- Invasion or postglacial invasion, 194, 366, 989, 1007 (see also Colonization)
- Isobar, 31, 32
- Isohyet, 33
- Isolation or Isolated, in Beringia or Yukon, 366, 372, 412, 415–416, 417, 476, 551, 552, 678, 779, 823, 824, 833, 835, 849, 989
 - correlated with wing loss, 978–979, 985
 - in East and West Beringia, 850
 - in peatlands, 679
 - on mountains, 10, 322–323, 370, 679, 783, 839, 997
 - to south, 813, 826, 835, 837, 839, 841
- Isotherm, 28, 29, 30 (see also Temperature)
- Isotope, see Oxygen isotope
- Lake or lakes, in Yukon, 14, 22, 24, 25, 26, 52, 896
 - and permafrost, 35 (see also Thermokarst)
 - as habitat for arthropods, 172, 173, 492, 615, 616, 791, 827, 835, 981
 - biological interactions, 859
 - species in, 164, 172, 220, 221, 307, 413, 516–519, 627, 632, 813, 837, 838, 849, 860, 982
- Lakes, access by, 226, 969
 - in past, 369, 860, 1006
 - fossils from, 917, 943, 944, 945, 947, 949
- Land bridge, see Bering land bridge
- Latitude, diversity and, see Diversity and latitude
- Latitude, Habitat and, see Habitats of insects, change with latitude
- Laurentide ice, ice-sheet or glacier, 9, 22, 23, 65, 455, 470, 790, 821, 822, 824, 830, 840, 848, 913, 917, 1000–1001, 1006
- Lentic habitats, availability or features of, in Yukon, 412, 413, 500, 791, 981, 1003, 1005
- Lentic species, occurrence in Greenland of, 828
 - occurrence in Yukon of, 164, 413–414, 493, 516–519, 827, 831, 858, 859, 981–982, 988
 - and freezing, 220, 858, 986
 - in temporary habitats, 307, 413 (see also Temporary or transient habitats)
 - relative to lotic species, 413, 518, 827, 854–856, 859, 988, 989
- Life cycle, 985–986, 1008
 - duration of, 118, 204, 220, 224, 639, 783, 833, 860, 985, 986, 1008 (see also Development, rate of)
 - of butterflies and moths, 783
 - of caddisflies, 833, 858, 860
 - of dragonflies, 195–196
 - of oribatid mites, 118
 - of orthopterans, 224
 - of robber flies, 639
 - of stoneflies, 204, 220
 - synchronized, 783, 985, 986
- Life span, of adult caddisflies, 828
 - of adult moths, 9, 987
- Life zone, 7, 8, 10, 191, 338, 344, 364, 408, 527, 870, 894, 965, 968, 980, 984, 997–999, 1008
 - and species zonation, 361, 362, 363, 515
- Limiting factors, in aquatic habitats, 221, 843, 858, 945, 982–983
 - in terrestrial habitats, 236, 896
- Limit of boreal fauna, 9, 10
- Limits, see Range limits
- Limnology, 25 (see also Lake)
- Litter, accumulation of, 984
 - as habitat, of ants, 904
 - of ground beetles, 981
 - of oribatid mites, 117, 118, 119, 142, 144
 - of spiders, 109, 110
 - with winglessness, 306
- Locality, see Collecting site
- Loess, 48, 63
 - and soil productivity, 66–67
- Lotic habitats, availability or features of, in Yukon, 173, 519, 856–859, 981, 1005
- Lotic species, occurrence in Yukon of, 164, 173, 307, 413, 493, 500, 515–519, 604, 827, 828, 858, 859, 981, 989, 1006
 - and freezing, 220, 858, 987
 - relative to lentic species, 518, 827, 854–858, 859, 988, 989
- Mammals and insects, 415, 604–608, 909, 948
- Mammal burrows or runs, species associated with, 664, 665, 675–677, 678, 981
- Mammalophily, in black flies, 568, 604–605, 607, 608

- Mammoth steppe or mammoth fauna, 60, 61, 62, 65, 68, 368, 477, 478, 916, 948
- Marine littoral habitats, mites in, 119, 142, 144, 146
- Marine oxygen isotope, see Oxygen isotope
- Marine sediments, 64
- Marsh, marshy, 172, 174, 307, 384, 413, 449, 464, 467, 479, 515–517, 518, 615, 791, 827, 848, 849, 853, 981
in past, 64, 479, 480, 860
- Mating, reduction of, 632, 979, 985
- Meadow, 63, 64, 449, 464, 466, 467, 480, 948
alpine meadow, 677, 678, 947
sedge meadow, 368, 370, 616, 1002
- Melanism, melanic, 849, 984, 985
- Metabolism, metabolic, 118, 857, 987
- Microenvironments or microhabitats, 982, 984
diversity of in Yukon, 980, 982, 1008
in Beringia, 480, 482
of ants, 903, 904
of ground beetles, 466
of oribatid mites, 119, 144, 146
of sphaerocerid flies, 674–677
- Microsites in vegetation, 43
- Migrants, in Yukon fauna, 970
- Migration, 162, 164, 236–238, 322, 323, 363, 917, 958, 1004 (see also Beringian emigrants; Beringian immigrants; Dispersal)
daily or seasonal, 220, 460
of trees, 68, 363
- Mimicry, 307, 308
- Models, for Beringian palaeoecosystems, 61, 64, 65
- Moisture, importance of for vegetation, current, 39, 44, 68, 980
in Pleistocene Beringia, 64, 65, 68, 1002
- Monophagy or monophagous, 302, 303–304, 378, 388, 401
- Mosaic, see Habitats
- Moss, in fossil peats, 63
- Moss, in Yukon habitats, 38, 41, 43, 44, 45, 48, 61, 500, 981, 1002
for ants, 904
for beetles, 414, 983, 1006
for caddisflies, 851, 861
for dragonflies, 169, 172, 174, 175, 982
for oribatid mites, 117, 119
for sphaerocerid flies, 674
for spiders, 109, 110
for true bugs, 302, 324
- Mountain or montane fauna or species, 8, 9, 145, 364, 365, 370, 372, 517, 518, 597, 604, 608, 655, 782, 827, 857, 909, 1006, 1008, 1009 (see also Alpine; Cordillera; Dispersal, barriers to; Refugia; Relict)
- Mountains, and habitats, 467, 597, 604, 776, 777, 827, 897, 980, 981, 989
in past, 64, 608, 953, 957, 1003
in Yukon, 13–23, 38–42, 53, 171, 791, 897, 966, 1009
- Mouthparts, of black flies, 563, 604–605, 608
- Movement or move, from Beringia, eastward or southward, 236, 830
into Beringia, northward or from south, 7, 8, 9, 10, 219, 236, 326, 679, 779, 854, 915, 1001, 1006 (see also Bering or Beringian land bridge...; Dispersal; Migration)
- Muskeg, 221, 823, 852, 860, 980
- Myrmecomorphy, in true bugs, 306, 307–308
- Nearctic, see especially Ranges
- Nest sites, of aculeates, 896–898
- New species, described, 487–488
- Northern Insect Survey, 245, 447, 569, 725
- Northernness, and Yukon occurrence, 975, 987–989, 1008
- Northwest Territories, fauna of, 117, 161, 163, 164, 192, 198, 203, 225, 235, 291, 301, 379, 401, 409, 419, 514, 518, 522, 541–543, 565, 587, 627, 631, 641, 691, 820, 870, 893, 975–976
- Number of species, see Composition; Diversity; Fauna
- Nunatak, 16, 22, 23, 301
- Nutrient-limited environments, 116
- Open forest, see Subarctic; Taiga
- Open or Open ground habitats or fauna, 48, 302, 411, 413, 415, 416, 460, 464, 466, 468, 469, 474, 475, 476, 479, 480, 481, 717, 718, 719, 778, 981, 1002, 1006
in past, 60, 63, 946, 949, 951, 1002
- Origins of the fauna, 9, 10, 954, 997, 1005, 1007 (see also especially Refugia; and see also Dispersal; Evolution)
in black flies, 606, 607, 608, 646
in caddisflies, 822, 824–825, 830 et seq.
in ground beetles, 446, 470, 471, 472, 475, 477
in leafhoppers, 363, 372, 373
in mayflies, 164
in robber flies, 657
in stoneflies, 218, 219
in weevils, 551–552
- Ornithophily, in black flies, 604–605, 607, 608
- Other Yukon species, see Additional species
- Overwintering, 118, 195, 196, 220, 306, 340, 639, 857, 858, 898, 986 (see also Cold hardness...; Diapause)
- Owls, as hosts for black flies, 569 et seq.
- Oxygen isotope, 59, 65
- Oxygen, resistance to low levels of, 858
- Pacific Coast, 54, 340, 360, 368, 517
- Palaeartic, see especially Range type; Ranges
- Palaeartic region, connections or affinities of Yukon species with, 8, 147, 679, 720, 916, 1008–1009 (see also Bering Strait..., not a separator...)
- Palaeoclimate, 61, 62, 64, 65, 68 (see also Climate)
- Palaeoecology, 23, 59–72, 369, 460, 482, 909 (see also especially Beringia, past habitats...)
- Palaeomagnetic reversals, 59
- Palynology, 60, 367, 480
- Pangaea, 146, 218, 235
- Pan trap, 3, 664, 678
- Paradox, Beringian, see Beringian
- Parasitoids, 869, 898–899, 984, 986, 988

- limitation of data on, 969, 987
 Parthenogenesis or parthenogenetic species, 118, 142, 146, 547, 606, 608, 833, 985
 Past conditions in Yukon, 59–72, 957–959, 1001–1002 (see also especially Beringia, past habitats...; see also under Arctic; Boreal; Dry; Grassland; and other habitats)
 Patterned ground, 36, 47
 Peat or peatland, fossils from, 63, 912, 941, 948, 955–956
 habitat, 36, 45–46, 47, 172, 173, 464, 492, 500, 616, 777, 983
 species associated with, 116, 119, 142, 172, 413, 464, 500, 515, 516, 517, 518, 519, 627, 675–677, 678, 679
 Permafrost, and fauna or their habitats, 117, 172, 860, 861, 896, 897, 957, 980, 981, 987
 and preservation of fossils, 914, 941
 and thermokarst, 944–945
 in the past, 68, 949, 957
 in Yukon, 35, 36, 45, 50, 52, 53, 343, 467, 616
 Phenology or phenological, 195, 984, 1008 (see also Emergence)
 Photographs, see under Habitat
 Photoperiod, 791, 859, 860, 986
 Physiography or physiographic, 9, 14–22, 50, 52, 897
 Phytolith, 63
 Phytophages, see Herbivores
 Plant associations, see Hostplant
 Pleistocene, see specific topics, e.g. Beringia...; Refugia
 Pliocene, conditions or changes in, 109, 917, 957–959, 1000, 1009
 connection, separation, or vicariance, 372, 821, 824, 830, 832, 836, 837, 839, 841, 842, 848, 852, 854, 917, 958, 959, 1000, 1004
 fauna of Yukon, reconstructed, 957–958
 fossils, see Fossils
 mammal dispersal, 823
 Pollen, see Fossils, pollen...
 Pollen influx, 60, 62–63
 Polygons, 38, 45, 47
 Polymorphism, see Wing polymorphism
 Polyphagy, 303–304, 308, 384
 Pond or pool (see also Rock pool; Saline habitats; Temporary or transient habitats)
 as habitat, 14, 22, 172, 174, 492, 615, 616, 791, 851, 859, 860, 861, 981, 982
 biological interactions in subarctic, 859
 collections from, 1, 3
 fossils in, 64, 912, 943, 956
 species in, 151, 169, 172, 174, 194, 195, 221, 306, 307, 413, 491, 500, 515, 516, 517, 519, 627, 787, 837, 838, 839, 843, 848, 851, 853, 860, 981–982, 983, 984, 986, 987, 1003
 alpine and subalpine, 518, 519
 Pool, see Pond
 Post-Pleistocene development of the fauna, 1006–1007
 Postglacial or post-Pleistocene, see specific topics, especially Dispersal; see also Colonization, etc.
 Precipitation, 18, 29–35, 65, 68, 492, 852, 896
 Predators, predation or predaceous, 221, 302, 303, 307, 308, 414, 416, 851–852, 859, 898–899, 968, 983–984, 988
 and habitat type, 413
 and range type, 318, 322, 323, 324, 326, 415
 relative occurrence in Yukon, 413, 856–857, 987, 989, 1008
 Production paradox, 61, 62, 66
 Radiation, of taxa, 234, 235, 236, 237
 Radiation, solar, 370, 869, 897, 915
 Rainfall, see Precipitation
 Range, and flight capacity, 468–469, 481, 546, 547–548, 552
 and habitat or zone, 107–110, 318, 322–326, 364–369, 415–416, 471 et seq., 515–517, 544–546, 550–551, 650, 652, 674–679, 776–782, 909, 997–999
 bicentric, 326, 470, 477, 482, 606, 1004
 multicentric, 1004
 unicentric, 325, 477
 within Yukon, see Distribution...
 zonal, 997–999
 Range limit or geographical limit, 191, 194, 199, 364, 384, 401, 517, 595, 627, 632, 651, 777, 778, 780, 781, 783, 897, 908, 909, 1005 (see also Ranges)
 Range types, 989–999, 1005–1006 (includes Biogeographic categories; Distributional categories, elements or patterns; Faunal elements; Geographical patterns; Range patterns; etc.—see also annotated lists of species in many chapters)
 definitions, 75, 173–176, 205, 247–248, 340, 344, 389, 408–409, 455–459, 493, 527–528, 640, 689–690, 820, 989–993
 in aculeate wasps, 895–896, 897
 in anthomyiid flies, 689–691, 692, 693, 715–721
 in ants, 907–909
 in aphids, 401–402
 in beetles, 410–412, 419–440
 in black flies, 566–567, 595–604, 606–608
 in butterflies and moths, 776–781
 in caddisflies, 820–854
 in chironomid midges, 628–632
 in delphacid planthoppers, 383–384
 in dragonflies, 173–176, 188–194, 197–198
 in ground beetles, 449–454, 455–459, 470 et seq.
 in leafhoppers, 361–369
 in mayflies, 162–164
 in oribatid mites, 122–129, 145–147
 in orthopterans, 235–238
 in robber flies, 640–641, 646–657
 in sphaerocerid flies, 674–679
 in spiders, 107–110
 in stoneflies, 218–219
 in true bugs, 308–326
 in water beetles, 494–499, 515–517
 in weevils, 550–552

[The following listings give detailed text references to particular types of ranges (see also the annotated lists for ranges of individual species), although many readers may prefer to use the general or group-by-group index entries above]

Ranges (see under Range types for terms equivalent to Ranges)

Amphiberingian, 247, 604, 820, 850–851 (see also Ranges, East-West Beringian)

Arctic or Arctic/Alpine, 75, 107, 108, 230, 343, 361, 362, 363, 365, 401, 402, 493, 495, 516, 517, 518, 628, 631, 693, 716–717, 718, 998, 999

Low Arctic, 361

Austral, 344, 361, 362, 363, 870, 894, 998

Beringian, 110, 191, 197, 198, 199, 362, 408, 409, 410, 449, 455–456, 469, 527–528, 550–552, 640, 646, 647, 649, 650, 653, 654, 656, 657, 674, 689, 715, 718, 720, 820, 824, 849–852, 859, 860, 979, 987, 989, 994, 995, 996, 999, 1002, 1004, 1005, 1009

East Beringian, 107, 108, 145, 147, 176, 205, 218, 219, 247, 320, 321, 340, 401, 402, 409, 410, 411, 419, 528, 544, 546, 547, 553, 566, 596, 597, 603, 604, 606, 607, 608, 640, 647, 649, 715, 778, 781, 782, 990, 991, 994, 995, 997–998, 1002, 1004, 1005, 1009

East-West Beringian, 107, 108, 109, 145, 147, 219, 247, 319, 320, 322, 324, 340, 364, 365, 366, 369, 372–373, 383, 410, 411, 419, 544, 546, 547, 715–716, 990, 991, 994, 995, 996, 1005

Beringian and glaciated western Nearctic, 493, 494, 516, 517

Boreal, 75, 108, 109, 110, 175, 188, 189, 192, 193, 364, 383, 389, 401, 402, 493, 496, 515–520, 639, 640, 646, 648, 653–656

Boreal/Temperate lowland, 716, 717

Low Boreal, High Boreal, 344, 345, 360, 361, 362, 363

Northern Boreal, 175, 188, 189, 190

Southern Boreal, 176, 189, 190

Western Boreal, 176, 189, 191, 389, 401, 402

Widespread Boreal, 175, 188

Boreal-Cordilleran, 716, 719

Boreomontane, 146, 189, 389, 401, 402, 640, 677, 679, 776, 780

Western boreomontane, 389, 402

Circumboreal, 247, 320, 322, 401, 402, 779, 838

Circumboreal Dune-dwelling, 718

Circumpolar, 108, 146, 320, 322, 340, 361, 383, 449, 455, 459, 469, 828, 990

Circumpolar Arctic/Alpine, 718

Circumpolar Arctic lowland, 718

Circumpolar High Boreal/Low Arctic, 718

Cordilleran, 107, 109, 162, 176, 190, 191, 193, 194, 493, 497, 516–518, 566, 596, 597, 602, 606, 607, 639, 640, 646, 648, 656, 657, 716, 717, 719, 993, 995, 996, 997, 1005, 1006, 1007

Cordilleran-Atlantic, 716

Cordilleran excluding Beringia, 247, 322, 324, 325

Cordilleran including Beringian, 247, 321, 325

Cordilleran-Prairie, 717, 719

Cordilleran-West Beringian, 107, 109, 994

Cosmopolitan, 146, 402, 717

Deciduous or Deciduous forest, 493, 499, 516, 517

Disjunct, or disjunction, 147, 162, 219, 243, 322, 325, 360, 364, 365, 384, 408, 527, 566, 597, 600, 657, 678, 679, 713, 716, 717, 725, 776, 778, 779, 780, 781, 782–783, 824, 827, 828, 829, 830, 831, 832, 833, 835, 836, 838, 846, 851, 896, 897, 969, 992, 993, 997, 998, 1004, 1005, 1006

East Palaeartic-Beringian, 716

East Palaeartic-Nearctic Boreal/Cordilleran, 717, 720

East Palaeartic-Nearctic Cordilleran, 718

Eastern Palaeartic and widespread western North American, 546

East Beringian, see Beringian

East-West Beringian, see Beringian

Grassland, 366, 373, 493, 495, 516, 517 (see also Grassland)

Holarctic or widespread Holarctic, 9, 107, 109, 110, 117, 146, 162, 163, 164, 173, 175, 188, 191, 205, 218, 219, 235, 236–237, 322, 340, 361, 362, 363, 372, 401, 402, 471, 493, 494, 495, 516, 520, 523, 543, 566, 595, 596, 604, 607, 640, 641, 648, 649, 655, 674, 675–676, 677, 678, 679, 721, 776–780, 820, 824, 825, 827–828, 830–839, 846, 851, 859, 860, 895, 897, 958, 989, 990, 994, 1005, 1008 (see also Ranges, Nearctic and Palaeartic)

Holarctic excluding Beringia or not in Beringia, 566, 596–597, 604, 820, 852–854

Holarctic including Beringian, 689, 717, 719, 720, 721

Montane (see also Ranges, Boreomontane)

Western Montane, 389, 401, 402, 408, 410, 411, 419, 527, 543, 544, 546, 776, 827 0

Nearctic, 107, 108, 109, 110, 146, 162, 173, 176, 199, 205, 218, 235, 237, 247, 321–322, 362, 363, 364, 365, 383, 401, 402, 410, 411, 523, 544, 566, 595–597, 598, 640, 647, 655, 675–676, 778, 780–781, 824, 828, 989, 994–999, 1005

East and West Nearctic, 676

Nearctic-Beringian or Nearctic including (East)

Beringian, 247, 321, 325, 449, 455, 456, 457, 469, 689, 691, 716, 719, 720, 780

Nearctic excluding (East) Beringia, 247, 321, 324, 325, 449, 455–459, 469, 689, 717, 719, 720, 990, 993, 994, 997

Northern Nearctic, 566, 597, 599, 604, 607, 990, 992, 1005–1006 (see also Ranges, East Beringian)

Northwestern Nearctic, 147, 205, 218, 676, 677

Transcontinental/Western Montane Nearctic, 410, 411, 419

Trans-Nearctic, 679

Western or West or Western Nearctic, 107, 109, 110, 176, 190, 191, 195, 205, 218, 363, 410, 411, 517, 527, 544, 550, 552, 566, 597, 601, 676, 677, 895, 993, 995, 996

Western Nearctic excluding Beringia, 247, 321, 324, 325, 326

Western Nearctic including Beringian, 247, 321, 325, 326

Widespread or Widespread Nearctic or North American, 176, 190, 196, 234, 238, 363, 364, 365, 389, 401, 402, 408, 410, 416, 526, 527, 550, 552, 553, 674, 679, 778, 780, 820, 825–830, 990, 992, 995, 996 (see also Ranges, Transcontinental)

Widespread Western, 410, 411, 419, 546

Nearctic-West Beringian, 107, 108, 109, 219, 383, 566, 596, 604, 607, 990, 994, 996, 1005

Nearctic and (plus) Palaeartic, 145, 543, 676, 677, 995

Widespread Nearctic and Palaeartic, 402, 779

Neotropical, 146, 218, 235, 322, 326

Other, 248, 322, 551, 720, 989

Palaeartic, 110, 117

Ranges (*continued*)

- Palaeartic-Beringian, 401, 402, 449, 455, 456, 469, 472, 1003
- Palaeartic-Cordilleran, 247, 320, 322, 323, 566, 596, 604, 607
- Palaeartic-Western Nearctic, 247, 321, 322, 323, 410, 411, 419
- Palaeartic-East Beringian, 107, 108, 109, 173, 175, 176, 190, 247, 319, 320, 322, 323, 324, 383, 401, 410, 470, 544, 550, 552, 566, 596, 604, 606, 607, 637, 640, 647, 649, 689, 690, 716, 719, 720, 721, 781, 820, 824, 839–849, 990, 991, 995, 996, 999, 1003, 1005
- Palaeartic-Transcontinental/Western montane Nearctic, 410, 411, 419
- Palaeartic-Transcontinental Nearctic, 410, 411, 419
- Palaeartic and Nearctic, 8, 109, 247, 320–321, 322, 383, 410, 411, 998, 999
- Palaeartic plus Nearctic Beringian, 780 (see also Ranges, Palaeartic-East Beringian)
- Palaeartic and Northwest Nearctic, 674
- Southern, 493, 498, 516, 517
- Subarctic or Subarctic/Alpine, 75, 108, 247, 320, 322, 326, 343, 345, 361, 362, 363, 365, 401, 402
- Transarctic, 322, 340
- Transboreal, 322
- Transberingian, 205, 218, 340
- Transcontinental, 9, 109, 162, 191, 205, 365, 410, 416, 493, 496, 498, 516, 517, 597, 827, 828, 830, 831, 832, 835, 837, 838, 853, 895, 896, 897, 907–908, 990, 1005 (see also Ranges, Nearctic, Widespread)
- Northern Transcontinental, 205, 218, 219, 340, 389, 408, 527, 543, 546, 544, 553
- Northern Transcontinental and Western Montane, 546
- Transition, 176, 190, 344, 360, 361, 362, 363, 364, 365, 998, 999
- Transpalaeartic-Beringian, 716
- Transpalaeartic-Nearctic Boreal/Cordilleran, 717
- Transpalaeartic-Nearctic Boreal/Low Arctic/Cordilleran, 718
- Transpalaeartic-Nearctic Cordilleran, 718
- Uncertain, 248, 455, 459
- Western Nearctic or Western, see Nearctic
- Widespread Nearctic or North American or Widespread, see Nearctic
- Widespread Palaeartic; Widespread North American, 546
- Widespread Palaeartic; Northern transcontinental and western montane North American, 546
- Widespread Palaeartic; Northern transcontinental North American, 546
- Widespread Palaeartic-East Beringian, 546
-
- Refugia or refugium, 9, 360, 369, 482, 822, 823, 1007
- Beringian or Yukon, 8, 23, 49, 54, 176, 192, 219, 220, 235, 236, 237, 238, 302, 322, 323, 325, 326, 363, 368, 369, 470 et seq., 477, 551, 553, 609, 640, 646, 679, 790, 821, 822, 824, 829, 832, 835, 838, 841, 848, 850, 851, 895, 897, 909, 987, 989, 1001, 1004, 1006, 1009
- Banks Island, 456
- coastal, 815, 854
- conditions in, see Beringia, past habitats...
- Cordilleran, 324
- disjunct or multiple, 833, 836, 1004 (see also Range, bicentric, multicentric; Ranges, disjunct)
- dispersal or spread from, 7, 823, 982, 1005
- eastern, 363
- importance of microenvironments in, 480–482, 482
- in Alaska, 366, 368, 479
- in mountains, 10, 323, 679, 825, 836
- Mackenzie River, 456
- northern, 325, 326, 897 (see also Refugia or refugium, Beringian)
- northwestern, 164, 219, 460, 471, 480 (see also Refugia or refugium, Beringian)
- Peary Land, 828
- southeastern, 219, 220, 679
- southern or continental, 9, 232, 322, 323, 324, 325, 326, 363, 460, 477, 482, 551, 552, 679, 790, 815, 825, 826, 827, 828, 830, 832, 837, 841, 987, 1004, 1005, 1006
- southwestern, 219, 220, 679
- western, 365
- Relict, geographic, 787, 824, 851, 861
- phylogenetic, 787, 824, 851, 861
- population, 225, 322, 323, 978, 1004
- vegetation, 61, 67, 366, 656, 781, 945, 947
- Relict distribution, 146
- Relict species, 366, 781, 852, 861, 1004
- Beringian, see especially Relict species, glacial
- glacial, 837, 838, 846, 849, 850, 851
- montane or mountain, 830, 838, 848–849, 978, 997, 1009
- Representation of taxa, see Composition
- Reproductive behaviour, 9, 987 (see also Mating)
- Resources, availability or use of, 164, 221, 414, 854–857, 858, 859, 982–984
- Restricted species, restricted or confined to Beringia, 9, 142, 147, 192, 366, 409, 410, 412, 415, 416, 441–444, 517, 544, 547, 548, 553, 652, 778, 779, 780, 784, 861, 994–997, 1002, 1005, 1009 (see also Endemic species)
- Richness, see Diversity
- Riparian species, 411, 413, 415, 455, 460, 467, 468, 469, 475, 476, 479, 480, 481, 550, 663, 675–677, 978, 979, 981
- River, Mackenzie, age of, 916–917
- River or rivers, access by, 226, 969
- and permafrost, 35
- as habitat for arthropods, 164, 173, 616, 791, 857, 858, 981
- dispersal by, 164, 220, 632, 1003
- fossils associated with, 60, 912, 917, 941, 944, 947, 949
- in Yukon, 14, 23–26, 154
- on Bering land bridge, 823, 832, 833
- species in, 173, 194, 413, 568, 608, 832, 835, 836, 853, 982 (see also Riparian; Valley)
- RNA, 326
- Roads or highways, 13, 16, 49, 206, 338

- access by, 50, 447, 568, 725
collections correlated with, 161, 226, 492, 594, 791, 969, 970, 1010
habitats along, 302, 466
Rock pool, 632, 979
Sage or sagebrush slopes or hillsides, 73, 110, 365, 656, 980 (see also *Artemisia*-grass)
Saline habitats, 9, 45, 46, 47, 307, 449, 464, 467, 516, 518, 519
Sampling, see Collecting
Sand dunes or Sandhills, see Dunes
Saprophages or Saprovores, 415, 983, 988, 1008
Scree, 9, 48, 119, 225, 325, 340, 726, 776–778, 780–782
Season length and variability, climatic, 30–35
Season length, response to, 195, 378, 783, 860, 985–987
Seasonal movement or migration, 221
Seasonality, of aquatic habitats, 492, 520, 981, 986 (see also Ice; Overwintering)
Sedge or sedges, 38, 41, 45, 52, 61, 65, 172, 173, 174, 175, 176, 324, 339, 368, 370, 464, 616, 674, 777, 778, 981, 1002 (see also Grasses and sedges)
Seepage habitats, 480, 615, 616, 981
Settlements, in Yukon, 14, 16
Sex ratio, 146, 979
Shredders, occurrence in Yukon, 220, 221, 856, 858, 982–983, 988
Shrubs, beetles associated with, 411, 449, 466–467, 469, 475, 476, 479, 480
Shrub tundra, 119, 147, 197 (see also Wet shrub-tundra)
Sibling species, 371, 565, 566, 569, 593, 595, 632, 824, 841
Sister groups, species or taxa, 10, 198, 234, 369, 370–372, 649, 653, 674–679, 812, 813, 816, 817, 824, 829, 830, 831, 832, 833, 835, 836, 837, 839, 841, 842, 843, 845, 846, 848, 850, 851, 958, 1004
Sites, for collecting, see Collecting
for fossils, see Fossils
Size, small, in Yukon, 194, 978, 984, 985, 987
Soil, and Pleistocene productivity, 66, 67
Soil, role of mites in, 116, 117, 119, 968, 982
Soil structure, 116
Soils in Yukon, 35–38, 41, 44, 966, 980
Southern steppe, see Steppe
South-facing slopes or hillsides, in Yukon, as habitat for arthropods, 48, 50, 302, 308, 325, 778, 869, 966, 980
collections from, 3, 4, 245
species of, 337, 361, 365, 369, 523, 656, 778, 780, 897, 1002, 1008
vegetation or conditions of, 38, 48, 54, 67, 338, 369, 656, 896, 897, 947
Speciation, 237, 412, 455, 475, 569, 824, 1009 (see also Evolution; Origin)
vicariant, 816, 830, 832, 837, 843, 848 (see also Vicariance)
Species complex, in beetles, 417, 475, 476, 477, 489
in black flies, 569, 595, 604
in caddisflies, 815, 816, 835, 853, 854
in other taxa, 288, 633, 665, 678, 903
(see also taxonomic notes in annotated lists)
Species, description of new, 487–488
Species-packing, 821, 840
Spring or springs, as habitat for arthropods, 9, 45, 480, 627, 981
species in, 220, 516, 519, 632, 675, 835, 857, 979, 981, 982
Steppe (including arctic steppe, southern steppe) habitat or species, of beetles, 413, 415, 416, 466, 471, 474, 481, 523, 544, 545, 549, 550, 551, 552
of butterflies and moths, 725, 777–783
of other taxa, 130, 142, 144, 146, 325, 326, 369, 652, 674–676, 678, 980, 998, 999
in past, 49, 60, 61, 62, 63, 65, 67, 68, 365, 368, 370, 372, 477, 479, 480, 552, 605, 823, 916, 947, 948, 1002
Stepping-stone dispersal, 1003
Storms, 28, 29, 32, 33, 34, 35
Stream or streams, and Cordilleran species, 604, 1006, 1009
as habitat of arthropods, 164, 173, 176, 194, 220, 221, 492, 597, 604, 616, 627, 791, 849, 850, 857, 858, 859, 981, 1005, 1008
colonization of, 857–858
fossils associated with, 947 (see also River)
in past, 823, 860, 1002, 1003
in Yukon, 14, 24
species of, 161, 164, 173, 176, 194, 217, 220, 221, 306, 500, 516, 519, 563, 568, 597, 608, 627, 813, 827, 831 et seq., 856, 857, 858, 859, 860, 982, 984
Structural or morphological adaptations, 9, 307, 632–633, 979
Subalpine, forest or vegetation, 38, 39, 43, 44–45, 48, 53, 54
species, 146, 365, 518, 519, 716, 717, 778
Subarctic, fossils or fossil sites, 915, 941, 955
habitats, 11, 14, 37, 43, 45, 53, 54, 343, 344, 966, 971, 980, 1009 (see also Taiga)
species, 8, 9, 10, 107–109, 117, 220, 221, 224, 237, 320, 322, 326, 361, 362, 363, 365, 372, 396, 401, 402, 470, 477, 830, 857, 859, 953, 985, 998, 999, 1005, 1006, 1008 (see also Ranges, Subarctic)
Subfossil, avoidance of term, 914
Subimagines, 153
Subspecies, multiple or several, 110, 236, 237, 372
Summer temperature, 26, 29, 31–34, 171, 172, 338, 340–344, 360, 896
Sun-warmed habitats, 361, 365, 369, 966, 980, 981, 985
Supercooling, 118, 858 (see also Cold hardiness...)
Swamp, 9
Swarming, 632
Sympatry or sympatric, 813, 836, 843, 846
Synanthropic species, 675–676, 678, 1007

- Synonymy, new, 117, 127, 134, 656, 812, 816, 817
- Systematic or taxonomic work, needs for, 117, 147, 288, 679, 780, 968, 1010
- Taiga, habitat or vegetation, 36, 39, 43, 45, 53
species of, 8, 471, 719, 776, 779, 909 (see also Subarctic)
- Talus, 17, 108
- Taphonomy, 941
- Taxonomic relationships, 412, 1004–1005 (see also Divergence; Sister groups...; Variation; Vicariance)
- Taxonomic work, see Systematic...
- Temperature, in Yukon, 26, 28–35, 171–172, 338, 344, 360, 896 (see also Climate, of Yukon; Growing season; Summer temperature)
- Temporary or transient habitats, 178, 306, 307, 378, 413, 518, 616, 787, 828, 848, 851, 854, 860, 861, 981, 986
vernal pools, 492
- Tephra, 67, 950
- Thelytoky, 118–119, 142, 985
- Thermokarst, 14, 944, 981
- Transcontinental, see Ranges
- Transient habitats, see Temporary...
- Treeline, 61, 164, 363, 628, 631, 687, 718, 780, 791, 828, 855, 897, 917, 982, 1004, 1009
change of or past location of, 945, 946, 953, 956
fossils near or representing, 10, 915, 918, 946, 951, 952
habitats or vegetation near, 44, 480
habitats or vegetation near, in past, 958 (see also Treeline, change of...)
importance or not as a boundary or range limit, 343, 361, 416, 627, 632, 853, 897, 980–981, 1003
in relation to past movements, 7, 9
species near, 10, 171, 172, 174, 175, 176, 197, 650, 830, 853, 909, 943, 945
- Trench, Tintina, 22, 23, 51, 53, 171, 247, 324, 396, 457, 458, 639, 792, 896, 989
- Shakwak, 22, 35, 43, 51, 161, 171, 173, 227, 639, 656, 792
- Trophic relationships or interactions, and correlation with Yukon occurrence, 856–858, 898–899, 982–984, 987–989
- Tundra and tundra vegetation, 38–41, 45, 52–54, 62, 63, 119, 146, 325, 467–468, 615, 616, 791, 855, 859, 860, 861, 980
in past, 49, 61–65, 67–68, 109, 324, 325, 326, 365, 369, 470, 471, 479, 480, 481, 553, 605, 608, 823, 825, 848, 851, 852, 853, 916, 917, 949, 951, 957, 959, 1000, 1001, 1002
age of, 825, 917, 959
- Tundra habitats, as represented by fossils, 915, 918, 943, 946, 951, 952, 954, 955, 956, 987
- Tundra species, 949, 998, 1002, 1003 (and see annotated lists for some taxa; see also Alpine; Arctic) and ranges, 8, 145, 146, 147, 321, 322, 324, 326, 365, 415, 416, 471, 550, 552, 652, 677, 678, 679, 718, 719, 778, 779, 780, 849, 1005, 1006, 1008
in tundra streams, 850, 857, 858, 860
of anthomyiid flies, 693, 719
of beetles, 411, 413, 416
of butterflies and moths, 723, 725, 777–782
of caddisflies, 787, 830, 837, 838, 839, 851, 860, 861
of ground beetles, 449, 445, 455, 468, 469, 474, 475, 476, 477, 480, 481
of leafhoppers, 361, 365, 366
of oribatid mites, 117, 119, 130, 142, 144, 145
of sphaerocerid flies, 663, 674–679
of stoneflies, 219, 221
of true bugs, 300, 301, 308, 323
of weevils, 523, 544, 545, 549
of other taxa, 108, 197, 627, 652, 897, 909
- Unglaciated areas, 14, 22, 23, 59, 145, 162, 219, 248, 321, 322, 323, 325, 366, 369, 409, 455, 469, 528, 715 et seq., 779, 821, 822, 896, 913, 916, 948, 965–966, 1001 (see also Refugia...; and see Ranges, Beringian)
- Unicentric, see Range, unicentric
- Univoltine, see Voltinism
- Valleys, influence on fauna, 161–162, 172, 194, 361, 364, 369, 518, 654, 656, 897, 966, 980, 981, 1006, 1007, 1008
in Yukon, 10, 17–23, 35, 54, 171, 174, 338, 366, 369, 457–458, 466, 896, 966
- Variation, morphological, or variable species, 978, 1004
in caddisflies, 812, 813, 815, 816, 820, 824, 829, 830, 832, 833, 835, 836, 837, 838, 839, 840, 846
in other taxa, 236, 476, 489, 527, 909
- Vegetation, types in Yukon, 38–45, 46, 48, 50, 52–54 (see also Habitats...; see also specific kinds of vegetation)
during Pleistocene, see especially Beringia, past habitats...
- Vertebrate fossils, see Fossils, vertebrate
- Vertebrates, see Birds; Mammals
- Vicariance or vicariant, 370–372, 654, 816, 824, 830, 831, 832, 837, 839, 841, 842, 843, 848, 850, 1004
- Vicars, 470, 476, 482, 606
- Voltinism or -voltine, 204, 220, 224, 306, 307, 340, 378, 837, 838, 839, 858, 860, 985–986, 987
- Warning colouration, see Aposematism
- Water (including sea and ocean), barrier or dispersal over, 632, 679, 721, 821, 823, 1003
- Watershed, 23, 24, 337, 369 (see also Drainage systems)
- Wet or moist tundra, 119, 615, 616 (see also Bog tundra)
in past, 823, 851, 852
species of, 144, 301, 322, 544, 545, 549, 627, 663, 674–678, 679, 693, 719, 981, 1002
- Wet shrub-tundra, 776–782 (see also Shrub tundra)
- Wetlands, 44, 45–46, 172–173, 464, 896 (see also Bog; Fen; Marsh; Peat or peatland; Swamp)
in past, 946
species of, 474, 475, 545, 546, 549, 550, 551, 980
- Wind, 31, 35, 361, 363, 365, 721, 777, 823, 979, 1006

- Wing dimorphism or polymorphism, 304–307, 308, 469, 474, 475, 476, 482, 550, 979
- Wing form, tabulated, 449–454, 528–529
- Wings, loss or reduction, Aptery, Brachyptery or Flightlessness, 978–979, 979, 985, 1008, 1009
- and habitat, 306, 340, 378, 416, 417, 468–469, 470, 474–477, 549, 784, 979, 985
- and latitude, 304
- and range, 326, 366, 415, 417, 469, 471, 476, 546, 552, 604, 1008
- and isolation, 978–979, 985
- and selective pressures or correlations, 304, 306–307, 468–469, 978–979
- in fossil record, 941, 952
- of beetles, 415, 416, 417
- of black flies, 604, 606, 609
- of butterflies and moths, 9, 723, 783–784, 987
- of caddisflies, 852
- of chironomid midges, 632
- of delphacid planthoppers, 377, 378, 384
- of ground beetles, 455, 460, 468–469, 470, 471, 475, 476, 481
- of leafhoppers, 340, 344, 365, 366, 371, 1003
- of sphaerocerid flies, 678
- of true bugs, 304–307, 308
- of weevils, 547–550, 552, 553
- Winter survival, see Overwintering
- Wisconsinan, see specific topics such as Beringia...;
Fossils; Glaciation; Glaciers
- Woodpeckers, feeding on ants, 903, 909
- Xeric habitats, 45, 48, 49, 302, 324, 325, 326, 479, 481, 517, 518, 550, 980 (see also Dry)
- Yukon project, 1–5, 7, 11, 152, 903
- Zonation, 361–363 (see also Range, zonal)
- Zoogeography, see especially Range types